Life in Dark Subsurface Oceans: Thermodynamic Constraints and Mixing Regimes

Leon Yilin Li[1]

Received January 9, 2026
Accepted May 7, 2026
Electronic access June 15, 2026

Background/Objective: Subsurface oceans on icy moons such as Europa, Enceladus, and Titan are among the most promising extraterrestrial habitats for life. Most habitability assessments focus on the presence of liquid water and chemical energy but do not quantitatively evaluate whether large-scale ocean mixing regimes permit the emergence of complex ecosystems. This paper investigates how the ratio of physical mixing timescales to biological evolution timescales may constrain the structure of hypothetical subsurface biospheres.
Methods: A heuristic dimensionless parameter, $\Pi_{mix} = \tau_{mix}/\tau_{bio}$, is constructed to compare vertical ocean mixing timescales to representative biological timescales. Order-of-magnitude estimates are computed for Earth and an idealized Europa ocean, with sensitivity analysis across plausible ranges of ocean depth, diffusivity, and biological timescale.
Results: For Earth, $\Pi_{mix} \approx 5 \times 10^{-3}$, indicating a well-mixed ocean on evolutionary timescales. For an idealized Europa ocean ($\sim$ 100 km deep), Π_{mix} ranges from $\sim$ 0.003 to $\sim$ 30 depending on assumed diffusivity, with plausible central estimates near or above unity, suggesting potentially slow mixing relative to biological change.
Conclusions: If Europa's ocean mixes slowly on biological timescales, any biosphere would likely be spatially restricted to localized energy sources rather than forming a globally extensive ecosystem. This heuristic framework, while speculative, offers a physically motivated approach for evaluating habitability beyond the simple presence of water and energy.

Keywords: astrobiology, Europa, subsurface ocean, ocean mixing, habitability, thermodynamics, extremophiles, icy moons

Introduction

Background and Context

For most of human history, discussions of extraterrestrial life assumed Earth-like planets bathed in starlight, with surface oceans and atmospheres similar to our own. However, over the past few decades, observations of icy moons in the outer Solar System have radically expanded our concept of habitable environments. Data from the Galileo and Cassini missions revealed strong evidence for global oceans beneath the frozen surfaces of Europa and Enceladus[1,2], while studies of Titan and other icy worlds suggest that subsurface liquid water may be common in the outer Solar System. At the same time, the discovery of thousands of exoplanets has raised the possibility that hidden oceans could exist in many planetary systems.

Problem Statement and Rationale

Current assessments of subsurface ocean habitability typically focus on two questions: does liquid water exist, and are chemical energy sources available? While these are necessary conditions, they are not sufficient for evaluating whether complex ecosystems—let alone intelligent life—could emerge. A critical factor that remains underexplored is the role of large-scale ocean mixing and transport in distributing energy, nutrients, and chemical species throughout the water column. On Earth, ocean circulation is fundamental to the global biosphere, transporting heat, dissolved gases, and nutrients across vast distances[3]. Without efficient mixing, even a chemically rich ocean might support only isolated, localized communities rather than a globally connected biosphere.

Existing thermal and geochemical models of icy moon oceans typically treat the ocean as a well-mixed reservoir or focus on local conditions near hydrothermal vents[4,5]. These approaches do not quantitatively address the timescale over which mixing redistributes energy relative to the timescale of biological evolution and ecosystem development. This gap motivates the present analysis.

Significance and Purpose

This paper introduces a heuristic dimensionless parameter, Π_{mix}, that compares the vertical mixing timescale of a subsur-

[1] *Lumiere Education*

face ocean to a representative biological timescale. By computing this ratio for Earth and an idealized Europa, we provide an order-of-magnitude framework for distinguishing ocean worlds where life might form globally connected ecosystems from those where life, if present, would be confined to localized energy oases. Although speculative, this framework complements existing habitability models by adding a transport dimension to the analysis.

Objectives

The objectives of this study are: (1) to review the thermodynamic and energy conditions necessary for life in dark subsurface oceans; (2) to construct a heuristic mixing parameter and apply it to Earth and Europa; (3) to assess the sensitivity of results to key assumptions; and (4) to discuss the ecological implications of different mixing regimes for hypothetical subsurface biospheres.

Scope and Limitations

This paper presents an order-of-magnitude, heuristic analysis rather than a detailed numerical ocean circulation model. The mixing parameter is an illustrative timescale ratio, not a formally derived similarity parameter. The analysis assumes simplified, one-dimensional ocean geometry and does not resolve lateral variations, convective plumes, or tidal forcing in detail. Biological timescales are represented by a single reference value with sensitivity analysis. The paper focuses on Europa as the primary case study but the framework is generalizable to other icy ocean worlds.

Astrobiology and Subsurface Oceans

The Drake Equation as a Framing Device

Astrobiology, the interdisciplinary study of life in the universe, seeks to understand where and how living systems might arise. A useful conceptual framework is the Drake Equation, which estimates the number of intelligent, communicative civilizations in our galaxy[6]:

$$N = R^* \times f_p \times n_e \times f_l \times f_i \times f_c \times L$$

where N is the number of communicative civilizations, R^* is the rate of star formation, f_p is the fraction of stars with planetary systems, n_e is the average number of habitable planets per system, f_l is the fraction of habitable planets where life arises, f_i is the fraction of biospheres developing intelligence, f_c is the fraction communicating, and L is the average civilization lifetime.

Important clarification: The Drake Equation is designed to estimate communicative civilizations, not microbial ecosystems. This paper uses the equation only as a brief historical entry point; the analysis that follows concerns habitability conditions for any form of life and does not attempt to constrain the higher-order Drake parameters (f_i, f_c). Modern exoplanet surveys[7,8] indicate that planetary systems are extremely common, with f_p approaching 1.0, though the fraction of truly habitable worlds remains poorly constrained due to detection biases favoring large, close-in planets.

Subsurface Habitable Environments

Earth's subsurface environments—aquifers, oil fields, and hydrothermal systems—harbor extensive microbial communities that can be partially or entirely independent of photosynthetically derived energy[9]. Subsurface biomass may equal or even exceed surface biomass[10]. These findings demonstrate that photosynthesis is not a prerequisite for life, opening the possibility that dark subsurface oceans could sustain biospheres driven entirely by chemosynthesis.

Jupiter's moon Europa has become a primary target for subsurface habitability studies. Galileo magnetometer data revealed an induced magnetic field consistent with a global, electrically conductive (likely salty) ocean beneath Europa's ice shell[11]. The ice shell is estimated at 15–25 km thick, with the ocean reaching depths of 60–150 km[1,12]. The development of a biosphere on Europa likely depends on two processes: the upward transport of reduced chemical species from water–rock interactions at the seafloor, and the downward delivery of oxidants from the irradiated surface through the ice shell[13,14]. Russell et al. (2017) propose that regions of high oxidant flux at the top of the Europan ocean may be zones of enhanced biological productivity[15].

However, recent research by Byrne et al. (2026) suggests that Europa's seafloor may be geologically quiet, with insufficient tidal stress to drive faulting, hydrothermal vents, or significant water–rock interaction under present conditions[16]. This finding, if confirmed, would substantially constrain the available chemical energy for any Europan biosphere and underscores the importance of understanding ocean mixing and transport as factors in habitability assessment.

Subsurface Oceans: Europa and Earth Compared

Earth's ocean habitability results from multiple factors: nutrient recycling through ocean circulation, pH buffering by the carbonate system, and moderate temperatures and pressures maintained by the atmosphere[3]. Ocean circulation transports heat, dissolved gases, and nutrients essential for life throughout the ocean at multiple spatial scales. Plate tectonics creates new oceanic crust, releasing critical nutrients such as phosphorus[17].

Europa's ocean differs fundamentally. It is isolated beneath a thick ice shell, lacks wind-driven surface mixing, and receives no sunlight. The primary energy sources are tidal heating from Jupiter's gravitational influence and geochemical reactions at the seafloor and ice–ocean interface[18,19]. Recent numerical ocean models suggest that Europa's ocean may nonetheless be dynamically active, with convection, eddies, and zonal jets driven by basal heating and the effects of ice shell melting and freezing on ocean salinity[20]. These models estimate horizontal eddy mixing coefficients of approximately $200 - 300\ m^2/s$, comparable to those estimated for Earth's Snowball Ocean period.

Thermodynamic and Energy Conditions for Life

Free Energy, Entropy, and Biological Systems

Living organisms maintain highly ordered internal states by continuously importing free energy from their environment and exporting entropy. This is fully consistent with the second law of thermodynamics: organisms are open systems, and the overall entropy of the organism plus its surroundings increases[21]. The thermodynamically relevant quantity for biological processes is Gibbs free energy (ΔG), not entropy directly. Metabolic reactions proceed when $\Delta G < 0$, coupling exergonic reactions (which release free energy) with endergonic reactions (which build biological order).

In the historical phrasing of Schrödinger[22], organisms "feed on negative entropy." While evocative, this language can be imprecise: organisms consume chemical potential energy stored in molecular bonds, not entropy per se. On potentially habitable worlds, hypothetical organisms would rely on available redox gradients as sources of free energy—for example, oxidant–reductant pairs in subsurface oceans or hydrothermal vent fluids. For Europa, candidate reactions include the oxidation of H_2, CH_4, and reduced sulfur species by O_2 or H_2O_2 delivered from the irradiated surface ice[23,24].

Energy Budgets for Europan Life

Thermodynamic constraints on Europan life can be bounded by estimating the available chemical energy flux. McCollom & Shock (1997) calculated that geochemical constraints on chemolithoautotrophic metabolism at seafloor hydrothermal systems yield energy fluxes on the order of $10^8 - 10^{10}$ J/year per vent field for reactions involving H_2, H_2S, and CH_4. Sahai et al. (2024) estimated that iron oxidation reactions ("iron snow") on Europa could potentially fuel microbial communities with power densities comparable to low-energy terrestrial chemosynthetic ecosystems.

These energy budgets place upper bounds on the total biomass that Europa's ocean could support. Even under optimistic assumptions, the total available chemical energy is orders of magnitude less than Earth's photosynthetically driven biosphere[25], suggesting that any Europan life would be energy-limited and spatially restricted to regions where redox disequilibria are maintained[4].

Linking Thermodynamics to the Mixing Analysis

In an astrobiological context, a living biosphere corresponds to an extremely small subset of all possible molecular configurations—a low-entropy macrostate maintained by persistent free-energy fluxes. The dimensional variables that describe the exchange of energy and entropy in a subsurface ocean (energy flux, temperature, pressure, characteristic length and time scales) naturally motivate the question: how rapidly does the ocean transport and redistribute the free energy required for metabolism? This question leads directly to the mixing analysis presented in the next section.

Dimensionless Mixing Analysis: A Heuristic Timescale Approach

Assumptions and Limitations

The following simplifying assumptions are adopted before presenting the analysis. The reader should evaluate all subsequent results in light of these limitations:

- **Diffusive approximation:** Vertical transport is represented by an effective vertical diffusivity κ. Advective processes, internal waves, and other non-diffusive transport mechanisms are absorbed into this single parameter.
- **Single characteristic ocean depth:** The ocean is approximated by a single representative depth H. For Earth, $H \approx 4 \times 10^3 m$. For Europa, $H \approx 1 \times 10^5 m$ ($\sim$ 100 km), consistent with typical interior models but not directly measured.
- **Single effective vertical diffusivity:** A constant κ is assumed throughout the water column. The baseline value $\kappa \approx 10^{-4} m^2/s$ is comparable to Munk's[26] global average for Earth's deep ocean, but directly observed open-ocean values are typically $\sim 10^{-5} m^2/s$, with values reaching 10^{-3}–$10^{-1} m^2/s$ near rough topography[27].
- **Representative biological timescale:** A single reference $\tau_{bio} = 10^6$ years characterizes long-term evolutionary or ecosystem-level adjustment, based on typical speciation timescales in the marine fossil record[28] and the duration of major diversification events[29]. Sensitivity to this choice is examined below.

- **One-dimensional, globally averaged ocean:** The ocean is treated as horizontally homogeneous. Lateral variations, regional circulation patterns, and localized structures (plumes, boundary currents) are not resolved.
- **Constant fluid properties and steady forcing:** Thermophysical properties and external forcing (tidal heating) are assumed constant.
- **Same τ_{bio} for Earth and Europa:** This isolates the effect of changing H and κ on Π_{mix} but implicitly assumes comparable evolutionary timescales, which may not hold.

Constructing the Mixing Parameter

To compare how rapidly an ocean mixes relative to the timescale of biological change, we define a dimensionless mixing parameter:

$$\Pi_{mix} = \tau_{mix}/\tau_{bio} = H^2/(\kappa \cdot \tau_{bio})$$

where H is ocean depth, κ is effective vertical eddy diffusivity, $\tau_{mix} \approx H^2/\kappa$ is the vertical mixing timescale, and τ_{bio} is a representative biological timescale.

Note: This parameter is a heuristic timescale ratio, not a formally derived Buckingham Pi group. A rigorous dimensional analysis would start from the governing equations for ocean convection (incorporating gravity, density, temperature contrast, rotation rate, and tidal forcing) and derive a minimal set of dimensionless groups. The present analysis uses only the timescale ratio as an illustrative comparison tool.

Earth as a Calibration Case

For Earth's ocean, adopting $H \approx 4 \times 10^3 m$ and $\kappa \approx 10^{-4} m^2/s$:

$$\tau_{mix} \approx H^2/\kappa = (4 \times 10^3)^2/10^{-4} \approx 1.6 \times 10^{11}\ s$$

With $\tau_{bio} = 10^6\ yr \approx 3.15 \times 10^{13}\ s$:

$$\Pi_{mix,Earth} \approx 1.6 \times 10^{11}/3.15 \times 10^{13} \approx 5 \times 10^{-3} \ll 1$$

Earth's ocean is well-mixed on the million-year biological timescale: vertical transport of heat and dissolved species is fast compared to slow evolutionary change.

Idealized Europa Case

For an idealized Europan ocean with $H \approx 1 \times 10^5 m$ and $\kappa \approx 10^{-4} m^2/s$:

$$\tau_{mix} \approx (10^5)^2/10^{-4} \approx 10^{14}\ s$$

$$\Pi_{mix,Europa} \approx 10^{14}/3.15 \times 10^{13} \approx 3$$

In this idealized case, Π_{mix} is of order unity or larger, suggesting that vertical mixing may be slow relative to biological timescales. However, this result depends critically on the assumed κ, which is highly uncertain for Europa.

Sensitivity Analysis

Because the assumed diffusivity directly determines whether Π_{mix} is much less than or greater than unity, a sensitivity analysis is essential. The table below shows Π_{mix} for Europa ($H = 100\ km$) across different values of κ and τ_{bio}:

$\kappa = 10^{-5} m^2/s$: $\tau_{bio} = 10^4\ yr \rightarrow \Pi \approx 3000$; $\tau_{bio} = 10^6\ yr \rightarrow \Pi \approx 30$; $\tau_{bio} = 10^8\ yr \rightarrow \Pi \approx 0.3$

$\kappa = 10^{-4} m^2/s$ **(baseline):** $\tau_{bio} = 10^4\ yr \rightarrow \Pi \approx 300$; $\tau_{bio} = 10^6\ yr \rightarrow \Pi \approx 3$; $\tau_{bio} = 10^8\ yr \rightarrow \Pi \approx 0.03$

$\kappa = 10^{-3} m^2/s$: $\tau_{bio} = 10^4\ yr \rightarrow \Pi \approx 30$; $\tau_{bio} = 10^6\ yr \rightarrow \Pi \approx 0.3$; $\tau_{bio} = 10^8\ yr \rightarrow \Pi \approx 0.003$

$\kappa = 10^{-1} m^2/s$: $\tau_{bio} = 10^4\ yr \rightarrow \Pi \approx 0.3$; $\tau_{bio} = 10^6\ yr \rightarrow \Pi \approx 0.003$; $\tau_{bio} = 10^8\ yr \rightarrow \Pi \approx 3 \times 10^{-5}$

Additionally, Ashkenazy et al. (2021) found that Europa's horizontal eddy mixing coefficient may be $\sim 200 - 300\ m^2/s$ based on numerical simulations. If horizontal mixing dominates transport, the effective mixing timescale would be much shorter than the purely vertical estimate, and Π_{mix} could be substantially less than unity even for deep oceans. This highlights the critical importance of ocean dynamics modeling—beyond the simple one-dimensional diffusive estimate used here—for evaluating Europan habitability.

The result is most sensitive to κ and H (which enter as H^2/κ), with τ_{bio} shifting the threshold linearly. Without observational constraints on Europa's actual mixing regime, the analysis cannot definitively place Europa in either the well-mixed or transport-limited regime.

Interpretation

$\Pi_{mix} \ll 1$ indicates an ocean that mixes many times within a biological timescale; nutrients and energy can be redistributed globally. $\Pi_{mix} \gtrsim 1$ indicates mixing comparable to or slower than biological change; energy and nutrients may remain concentrated near their sources. The transition is gradual, not a sharp boundary: values of $\Pi_{mix} \sim 0.1 - 1.0$ represent an intermediate regime where mixing provides partial but incomplete redistribution. Characterizing the precise boundaries of these regimes would require detailed numerical circulation models (e.g., Ashkenazy et al. 2021; Lemasquerier et al. 2023[30]) that resolve convection, stratification, and tidal forcing.

Potential Biological Adaptations in a Transport-Limited Ocean

Extremophile Analogues

Extremophiles are organisms that thrive in extreme environments such as hydrothermal vents, hypersaline brines, and high-pressure deep-sea settings[31]. On Earth, these organisms define the known limits of life in terms of temperature, pressure, pH, and energy availability. In the context of icy ocean worlds, deep-sea vent communities and other marine extremophiles serve as empirical models for chemotrophic ecosystems that might arise in dark, high-pressure subsurface oceans powered by redox disequilibria rather than sunlight.

Predicted Metabolisms and Adaptations

If Europa's ocean is in the slow-mixing regime ($\Pi_{mix} \gtrsim 1$), energy and nutrients would remain concentrated near their sources rather than being redistributed globally. The most plausible metabolisms would be chemolithoautotrophic ones exploiting local redox gradients—for example, oxidizing H_2, CH_4, or reduced sulfur and iron species released by seafloor water–rock interactions, using CO_2 or sulfate as electron acceptors[4,24,32]. Oxidants such as O_2 and H_2O_2 from Europa's irradiated surface ice could further enhance local redox disequilibria at the ice–ocean interface[13,15].

Organisms in such environments would likely be psychrophilic (cold-adapted) and barophilic (pressure-adapted), with flexible enzymes, pressure-stable membranes, and stress-response systems allowing metabolism near the freezing point and at high pressures, analogous to Earth's polar and hadal microorganisms[33,34].

Ecosystem Structure in a Patchy Environment

Independent of the Π_{mix} analysis, the spatial distribution of energy sources on Europa—hydrothermal vent fields, fracture zones, ice–ocean interfaces—would naturally create a patchy habitat structure. On Earth, hydrothermal vent communities are characteristically patchy, with dense microbial mats and macrofauna concentrated around active vents and sparse life in the intervening deep ocean[35,36]. If similar patchiness exists on Europa, the biosphere would consist of discrete oases rather than a continuous, globally connected ecosystem.

Between these patches, slow mixing and weak gradients would favor ultra-low-energy survival strategies such as dormancy, extremely slow growth, and opportunistic dispersal, paralleling deep-sea and subsurface microbes on Earth that persist in energy-poor sediments[37,38].

From Microbial Ecosystems to Complexity

Energy Constraints and Major Transitions

Earth's history demonstrates that the transition from microbial ecosystems to complex life is not automatic but depends strongly on energy availability and key evolutionary transitions. For over three billion years, life on Earth was almost entirely microbial; complex animals appeared only after the evolution of eukaryotic cells with mitochondria and substantial increases in oxygen and metabolic power per cell[39,40]. The rise of multicellular animals and large brains is tightly coupled to transitions that increased available metabolic power per unit biomass[41,42].

Chemosynthetic ecosystems on Earth demonstrate both the potential and the constraints of energy-limited biospheres. Total chemical energy flux and resulting global biomass at deep-sea hydrothermal vents are small compared to that supported by photosynthesis[23,25,43]. If Europa's ocean is powered solely by seafloor reactions and ice-delivered oxidants, the total energy budget would constrain the biosphere to modest global biomass and high spatial heterogeneity.

Ecological Complexity Within Chemosynthetic Oases

Within localized energy oases, however, microbial communities could achieve ecological complexity. On Earth, chemoautotrophic vents host intricate consortia in which bacteria and archaea partition redox niches, form syntrophic partnerships, and support multi-trophic food webs including worms, crustaceans, and mollusks (Nakamura & Takai 2014, 2015[44,45]; Zeng et al. 2021). Europan analogues might involve biofilms and microbial mats around vent chimneys or fracture networks, with grazing and predatory microbes, viruses, and possibly small multicellular consumers evolving wherever energy fluxes and stability remain high over geological timescales.

On the Question of Intelligence

Speculative note: On Earth, advanced cognition has evolved independently in corvids, primates, and cephalopods, demonstrating that intelligence is not restricted to one lineage or body plan[46,47]. Cephalopods in particular show that large brains and sophisticated problem-solving can arise in fully marine, three-dimensional environments. However, we emphasize that the present analysis provides no evidence for or against the evolution of intelligence on Europa. Demonstrating a causal link between slow ocean mixing and intelligence evolution would require quantitative models of standing biomass, population sizes, encounter rates, predator–prey dynamics, and cost-benefit analyses of nervous systems—none of which are provided here. The most realistic expectation is a biosphere rich in microbial and perhaps small multicellular diversity, with the evolution of large, intelligent organisms being possible only under rare, locally favorable energy conditions.

Potential Sensory Adaptations in Dark Oceans

In the complete absence of sunlight, sensory evolution in any Europan biosphere would be biased away from vision and toward alternative modalities. By analogy with Earth's deep-sea organisms, the most likely sensory systems would include: chemoreception (detecting dissolved chemicals and gradients), mechanoreception (sensing pressure waves, vibrations, and currents), and possibly electroreception (detecting weak electric fields generated by biological or geochemical processes). These modalities could enable organisms to construct detailed internal maps of their environment and communicate through sound, chemicals, or electric fields, as many deep-sea organisms do on Earth.

Conclusion

Key Findings

This paper introduces a heuristic dimensionless mixing parameter, Π_{mix}, to compare ocean mixing timescales with biological evolution timescales on icy ocean worlds. For Earth, $\Pi_{mix} \approx 5 \times 10^{-3}$, confirming that our ocean is well-mixed on evolutionary timescales. For an idealized Europa ocean, Π_{mix} ranges from $\sim 3 \times 10^{-5}$ to ~ 3000 depending on assumed parameters, with central estimates near or above unity, suggesting that vertical mixing may be slow relative to biological change.

Implications

If Europa's ocean is in the transport-limited regime, any biosphere would most likely consist of psychrophilic, barophilic chemolithoautotrophic microbes adapted to narrow, energy-rich niches near hydrothermal vents, fracture zones, or ice–ocean interfaces. These communities could be ecologically complex but globally sparse. Earth's history indicates that the evolution of large, complex organisms requires sustained high power per cell, which Europa's energy-limited environment may not broadly provide.

Limitations

The analysis has several important limitations. The mixing parameter is a heuristic timescale ratio, not a formally derived dimensionless group. Results depend critically on the assumed eddy diffusivity, which is essentially unknown for Europa. The one-dimensional, diffusive approximation neglects potentially important three-dimensional dynamics including convection, tidal forcing, and zonal jets. The biological timescale is Earth-centric and may not apply to extraterrestrial biospheres. The recent Byrne et al. (2026) finding of a potentially inactive Europan seafloor, if confirmed, would fundamentally alter the energy source assumptions underlying this analysis.

Recommendations for Future Work

Future studies should: (1) couple the mixing timescale framework with detailed numerical ocean circulation models (e.g., Ashkenazy et al. 2021) to obtain physically grounded estimates of transport in icy moon oceans; (2) compute free-energy yields for specific Europan redox couples and bound the resulting biomass; (3) develop quantitative models linking mixing regimes to ecosystem structure, including standing biomass, dispersal distances, and ecological connectivity; and (4) use data from the Europa Clipper mission (arriving 2031) to constrain ice shell thickness, ocean depth, and mixing processes.

Closing Thought

The search for life in dark subsurface oceans challenges us to think beyond the Earth-centric assumption of well-mixed, sunlit oceans. The framework presented here, while preliminary, suggests that the transport regime of an ocean may be as important as the presence of liquid water and chemical energy in determining whether life can flourish. As we prepare to explore these hidden worlds with missions like Europa Clipper, understanding ocean dynamics will be essential to interpreting whatever we find—or do not find—beneath the ice.

Acknowledgments

I thank mentor Adrian Lopez (Harvard University) for his guidance in the development of this research project.

References

1 R. Pappalardo, J. Head, R. Greeley, R. Sullivan, R. Lorenz, S. Giese, R. Greenberg, P. Helfenstein, D. Senske and G. Hoppa, Does Europa have a subsurface ocean? Evaluation of the geological evidence. Journal of Geophysical Research: Planets, 104, 24015–24055, 1999. https://doi.org/10.1029/1998JE000628.

2 F. Postberg, N. Khawaja and B. Abel, Macromolecular organic compounds from the depths of Enceladus. Nature, 558, 564–568, 2018. https://doi.org/10.1038/s41586-018-0246-4.

3 C. Langmuir and W. Broecker, How to build a habitable planet. Revised edition, Princeton University Press, 2012.

4 K. Hand, C. Chyba, J. Priscu, R. Carlson and K. Nealson, Astrobiology and the potential for life on Europa. In: Europa (Eds. Pappalardo,

R.T., McKinnon, W.B., and Khurana, K.K.), University of Arizona Press, 589–630, 2009.

5 S. Vance, M. Panning and S. Stähler, Geophysical investigations of habitability in ice-covered ocean worlds. Journal of Geophysical Research: Planets, 123, 180–205, 2018. https://doi.org/10.1002/2017JE005341.

6 F. Drake, S. Shklovskii and C. Sagan, Discussion at Space Science Board-National Academy of Sciences Conference on Extraterrestrial Intelligent Life. Green Bank, WV, Holden-Day, 1961-11. As described in: I.

7 A. Cassan, D. Kubas and J.-P. Beaulieu, One or more bound planets per Milky Way star from microlensing observations. Nature, 481, 167–169, 2012. https://doi.org/10.1038/nature10684.

8 C. Dressing and D. Charbonneau, The occurrence of potentially habitable planets orbiting M dwarfs estimated from the full Kepler dataset and an empirical measurement of the detection sensitivity. The Astrophysical Journal, 807, 45, 2015. https://doi.org/10.1088/0004-637X/807/1/45.

9 J. Fredrickson and T. Onstott, Microbes deep inside the Earth. Scientific American, 275, 68–73, 1996.

10 T. Hoehler and L. Poole, Habitability, life, and the biosphere. Oxford Research Encyclopedia of Planetary Science. Oxford University Press, 2018.

11 M. Kivelson, K. Khurana, C. Russell, M. Volwerk, R. Walker and C. Zimmer, Galileo magnetometer measurements: a stronger case for a subsurface ocean at Europa. Science, 289, 1340–1343, 2000. https://doi.org/10.1126/science.289.5483.1340.

12 J. Anderson, G. Schubert, R. Jacobson, E. Lau, W. Moore and W. Sjogren, Europa's differentiated internal structure: inferences from four Galileo encounters. Science, 281, 2019–2022, 1998. https://doi.org/10.1126/science.281.5385.2019.

13 R. Carlson, M. Anderson and R. Johnson, Hydrogen peroxide on the surface of Europa. Science, 283, 2062–2064, 1999. https://doi.org/10.1126/science.283.5410.2062.

14 B. Schmidt, The astrobiology of Europa and the Jovian system. In: Planetary Astrobiology (Eds. Meadows, V.S., Arney, G.N., Schmidt, B.E., and Marais, D.J.), University of Arizona Press, 185–216, 2020.

15 M. Russell, A. Murray and K. Hand, The possible emergence of life and differentiation of a shallow biosphere on irradiated icy worlds. Astrobiology, 17, 1265–1273, 2017. https://doi.org/10.1089/ast.2016.1600.

16 P. Byrne, H. Dawson, C. Klimczak, L. Jozwiak and S. Kattenhorn, Little to no active faulting likely at Europa's seafloor today. Nature Communications, 17, 2026. https://doi.org/10.1038/s41467-025-67151-3.

17 J. Korenaga, Plate tectonics and planetary habitability: current status and future challenges. Annals of the New York Academy of Sciences, 1260, 87–94, 2012. https://doi.org/10.1111/j.1749-6632.2011.06276.x.

18 C. Chyba and C. Phillips, Possible ecosystems and the search for life on Europa. Proceedings of the National Academy of Sciences, 98, 801–804, 2001. https://doi.org/10.1073/pnas.98.3.801.

19 M. Běhounkova, G. Tobie, G. Choblet and O. Čadek, Tidally induced magmatic pulses on the oceanic floor of Jupiter's moon Europa. Geophysical Research Letters, 48, e2020GL090077, 2021. https://doi.org/10.1029/2020GL090077.

20 Y. Ashkenazy, E. Tziperman and J. Chiang, Dynamic Europa ocean shows transient Taylor columns and convection driven by ice melting and salinity. Nature Communications, 12, pg. 6376, 2021. https://doi.org/10.1038/s41467-021-26710-0.

21 B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts and P. Walter, Molecular biology of the cell. 5th edition, Garland Science, 2008.

22 E. Schrödinger, What is life? The physical aspect of the living cell. Cambridge University Press, 1944.

23 T. McCollom and E. Shock, Geochemical constraints on chemolithoautotrophic metabolism by microorganisms in seafloor hydrothermal systems. Geochimica et Cosmochimica Acta, 61, 4375–4391, 1997. https://doi.org/10.1016/S0016-7037(97)00241-X.

24 N. Sahai, D. LaRowe and J. Senko, Bioenergetics of iron snow fueling life on Europa. Proceedings of the National Academy of Sciences, 121, pg. e2316452121, 2024. https://doi.org/10.1073/pnas.2316452121.

25 Y. Bar-On, R. Phillips and R. Milo, The biomass distribution on Earth. Proceedings of the National Academy of Sciences, 115, 6506–6511, 2018. https://doi.org/10.1073/pnas.1711842115.

26 W. Munk, Abyssal recipes. Deep-Sea Research, 13, 707–730, 1966.

27 K. Polzin, J. Toole, J. Ledwell and R. Schmitt, Spatial variability of turbulent mixing in the abyssal ocean. Science, 276, 93–96, 1997. https://doi.org/10.1126/science.276.5309.93.

28 D. Rabosky, Speciation rate and the diversity of fishes in freshwaters and the oceans. Journal of Biogeography, 47, 1207–1217, 2020. https://doi.org/10.1111/jbi.13839.

29 D. Erwin, M. Laflamme, S. Tweedt, E. Sperling, D. Pisani and K. Peterson, The Cambrian conundrum: early divergence and later ecological success in the early history of animals. Science, 334, 1091–1097, 2011. https://doi.org/10.1126/science.1206375.

30 D. Lemasquerier, A. Grannan, J. Vidal, D. Cebron, B. Favier, M. Bars and J. Aurnou, Europa's ocean translates interior tidal heating patterns to the ice-ocean boundary. AGU Advances, 4, pg. e2023AV000994, 2023. https://doi.org/10.1029/2023AV000994.

31 L. Rothschild and R. Mancinelli, Life in extreme environments. Nature, 409, 1092–1101, 2001. https://doi.org/10.1038/35059215.

32 J. Weber, T. Marlin, M. Prakash, B. Teece, K. Dzurilla and L. Barge, A review on hypothesized metabolic pathways on Europa and Enceladus. Life, 13, pg. 1726, 2023. https://doi.org/10.3390/life13081726.

33 P. Rampelotto, Extremophiles and extreme environments. Life, 3, 482–485, 2013. https://doi.org/10.3390/life3030482.

34 N. Merino, H. Aronson and D. Bojanova, Living at the extremes: extremophiles and the limits of life in a planetary context. Frontiers in Microbiology, 10, 780, 2019. https://doi.org/10.3389/fmicb.2019.00780.

35 L. Levin, A. Baco and D. Bowden, Hydrothermal vents and methane seeps: rethinking the sphere of influence. Frontiers in Marine Science, 3, pg. 72, 2016. https://doi.org/10.3389/fmars.2016.00072.

36 L. Mullineaux, Deep-sea hydrothermal vent communities. In: Marine community ecology and conservation, Sinauer Associates, 382–404, 2014.

37 X. Zeng, K. Alain and Z. Shao, Microorganisms from deep-sea hydrothermal vents. Marine Life Science and Technology, 3, 204–230, 2021. https://doi.org/10.1007/s42995-020-00086-4.

38 P. Higgins, Modelling extraterrestrial habitability, biomass and biosignatures in icy world oceans. PhD thesis, University of Edinburgh, 2022.

39 N. Lane and W. Martin, The energetics of genome complexity. Nature, 467, 929–934, 2010. https://doi.org/10.1038/nature09486.

40 E. Szathmáry and J. Smith, The major evolutionary transitions. Nature, 374, 227–232, 1995. https://doi.org/10.1038/374227a0.

41 C. Reinhard, N. Planavsky, S. Olson, T. Lyons and D. Erwin, Earth's oxygen cycle and the evolution of animal life. Proceedings of the National Academy of Sciences, 113, 8933–8938, 2016. https://doi.org/10.1073/pnas.1521544113.

42 N. Lane, How energy flow shapes cell evolution. Current Biology, 30, 471–476, 2020. https://doi.org/10.1016/j.cub.2020.03.055.

43 J. McNichol, S. Sylva, F. Thomas, C. Taylor, S. Sievert and J. Seewald, Primary productivity below the seafloor at deep-sea hot springs. Pro-

ceedings of the National Academy of Sciences, 115, 6756–6761, 2018. https://doi.org/10.1073/pnas.1804351115.

44 K. Nakamura and K. Takai, Theoretical constraints of physical and chemical properties of hydrothermal fluids on variations in chemolithotrophic microbial communities. Progress in Earth and Planetary Science, 1, pg. 5, 2014. https://doi.org/10.1186/2197-4284-1-5.

45 K. Nakamura and K. Takai, Geochemical constraints on potential biomass sustained by subseafloor water-rock interactions. In: Subseafloor Biosphere Linked to Global Mantle Dynamics (Eds. Ishibashi, J., Okino, K., and Sunamura, M.), Springer, 15–29, 2015.

46 P. Amodio, M. Boeckle, A. Schnell, L. Ostojic, G. Fiorito and N. Clayton, Grow smart and die young: why did cephalopods evolve intelligence?. Trends in Ecology and Evolution, 34, 45–56, 2019. https://doi.org/10.1016/j.tree.2019.05.005.

47 G. Roth, Convergent evolution of complex brains and high intelligence. Philosophical Transactions of the Royal Society B, 370, pg. 20150049, 2015. https://doi.org/10.1098/rstb.2015.0049.

