

Why the Klein Bottle Cannot Enclose Volume in R^3 : An Expository Topological Analysis

Zeyu Wang¹

Received April 17, 2026

Accepted August 19, 2026

Electronic access October 15, 2026

The Klein bottle is a classical non-orientable surface in topology that cannot be embedded in three-dimensional Euclidean space without self-intersection. This paper presents an expository analysis explaining why the Klein bottle cannot enclose a volume in R^3 . We examine orientability, embeddings and immersions. Using topological arguments and properties of smooth surfaces, we show that any surface enclosing a volume R^3 must be orientable. We further discuss why the Klein bottle may be immersed in R^3 with self-intersection but also an embedding in R^4 . Results show the relationship between topology and geometric constraints. The non-orientability of the Klein bottle and its inability to embed in R^3 are known; however, these topics are often presented separately and briefly in textbooks. This paper combined these concepts into a single logical framework, demonstrating how orientability, embeddings, immersions, and classical theorems collectively explain why the Klein bottle cannot enclose volume in R^3 .

Keywords: orientability, embedding, Euclidean space, boundaries, immersed.

Introduction

The Klein bottle was found in 1882 by German mathematician Felix Klein, and it is considered to be a classic example in topology because of its non-orientability and unusual geometric properties. Unlike familiar surfaces such as spheres and tori, the Klein bottle cannot be present in three-dimensional Euclidean space without self-intersection. Modern topology literature frequently uses the Klein bottle as an example of quotient constructions and non-orientable surfaces.^{1,2} Differential topology textbooks further use the Klein bottle as an example to illustrate quotient manifolds, smooth structure, and immersion versus embedding.³

Orientability is a fundamental property of surfaces and determines where consistent orientation may be defined across an entire manifold. A surface is considered to be orientable if it can be assigned continuous normal vectors globally.³ Classical examples of non-orientable surfaces are the Möbius strip and the Klein bottle. Non-orientable surfaces possess the property that the normal vector and orientation reverse, which creates important geometric restrictions and influences the application of some mathematical theorems.

For example, a sphere S^2 , defined by

$$x^2 + y^2 + z^2 = r^2$$

Which encloses a volume,

$$\text{Vol}(V) = \frac{4}{3} \pi r^3$$




Figure. 1 A sphere with outward-pointing unit normal vectors

Such a surface is orientable, meaning it has a continuous

¹ Balmoral hall high school, Canada

unit normal vector field $\vec{n}(p)$ over all points $S \ni p$. As Figure 1 shows, all the unit normal vectors are pointing outward over all points.




Figure. 2 Immersion of Klein bottle in R^3

As Figure 2 shows, the Klein Bottle has unit vectors pointing outward when on the surface but pointing inward when approaching the “opening” of the Klein Bottle. At first glance, the Klein Bottle appears to be a closed surface that has a bounded volume. However, Klein Bottle fails to separate the space inside and outside.

The distinction between embeddings and immersions is important in understanding the Klein bottle.⁴ Both immersion and embeddings require the derivative matrix to have maximal rank equal to the dimension of the starting space M . The difference is that an immersion only requires the derivative of a map to have constant maximal rank locally, meaning that it can be an “embedding” around a certain point, while an embedding holds a stricter topological structure in that the map needs to be homomorphic onto its image too, meaning that it

is an “immersion” without self-intersections.^{3,5} Existing literature proved that the Klein bottle can be immersed into R^3 but does not embed in R^3 . Self-intersection in three-dimensional space occurred due to the dimensional limitations and the non-orientable property of the Klein bottle.

The Möbius strip and the Klein bottle are often talked about together as fundamental examples of non-orientable 2-manifolds. The Klein bottle is constructed by pasting two Möbius strips together along their boundary, physically by splitting a square with mixed orientations (reserving one pair) along a diagonal and flipping one of the triangular pieces over. After pasting the sides with reserved orientations, a Klein bottle was formed. Topologically, a Klein bottle is equivalent to the connect sum of two real projective planes $RP^2 \# RP^2$.

In the last century, the related work on Klein bottle mainly focused on its classification of non-orientable surfaces and manifolds. Subsequent research has studied its algebraic-topological properties, including its fundamental group and topological complexity, which study found out the fundamental group of the Klein bottle is defined as $\pi_1(K) = \langle a, b \mid abab^{-1} = 1 \rangle$ and the topological complexity $TC(K)$ of the Klein bottle K is 4.^{6,7} Furthermore, in algebraic topology, researchers use different approaches to the generalization of the Klein bottle, extending it to n dimensions, as well as computing its integral cohomology algebra. The topological properties of the Klein bottle K_n varied on n , specifically its parity properties. K immerses into Euclidean space R^{n+1} and embeds into R^{n+2} without self-intersection.⁸

Later, mathematicians further studied the immersions of the Klein bottle, indicating that a simplified version requires exactly nine vertices to immerse the Klein bottle in R^3 . From Euler’s formula that any standard triangulations of a Klein bottle must have at least eight vertices, counterintuitively the theoretical eight vertices aren’t enough due to the geometric constraints in R^3 .⁹ Adding another vertex provides an extra dimension to move; just like crossing a small stream, jumping into the “third dimension” helps to avoid “crashing” into the stream.

Likewise, the Klein bottle is used not only as a main study subject but also as a fixed topological base for problems in geometry. In spectral geometry, mathematicians maximized the first eigenvalue on a Klein bottle, which explicitly shows the effect of non-orientability on geometric shapes. Meanwhile, the first eigenfunctions of the Klein bottle metric admit a minimal isometric embedding into a four-dimensional sphere S^4 .¹⁰ Other studies in differential geometry analyze a specific non-orientable minimal surface in Lawson’s bipolar surface $\tau_{1,1}$. Whose surface is a Klein bottle.¹¹

Recent studies have explored the stability of systolic inequalities that provide an alternative proof for Bavard’s optimal lower bound on the systolic area of a Klein bottle and further extend to the Möbius strip. By the estimation on the sys-

tolic defect $\alpha_{\text{sys}} - \frac{2\sqrt{2}}{\pi}$ using the L^2 -distance of the conformal factor from the systolic area minimizing metric. Similar estimates for metric also appear on the Möbius strip.¹² Another recent published paper proposed the idea that Klein bottle cosmology is a theoretical physics framework, physicists and mathematicians together explore a higher-dimensional universe constructed with the product of Minkowski space and Klein bottle. A huge step towards our universe's evolution, the topology of the universe itself potentially causes the creation of matter-antimatter asymmetry.¹³

Methodology

This paper used keywords such as “Klein bottle,” “topology,” and “Euclidean space” to search for relevant literature. In terms of enhancing the precision, literature and sources are selected from verified online journals and textbooks on reliable publication sites like arXiv and Cambridge University Press within the time range of 1926 to present. Since this study is based on existing literature and topology textbooks, sources are qualified if and only if they are cited multiple times by other researchers or are being used as teaching resources in universities or colleges, while authors must publish multiple research papers in their careers for credibility and reliability purposes, and their key findings and reinforcing reports must contribute to future studies and help develop mathematics over time.

First, introduce some key theorems and definitions to set up the base of this study, then build upon it by proving special properties of the Klein bottle, for example, non-orientability, embedding, and immersion. Ultimately, linked to the divergence theorem, the paper presented an expository topological analysis of the Klein bottle.

Theoretical Framework

1. Theorem (Jordan-Brouwer separation theorem)¹⁴ Let $X \subset R^n$ be a compact connected hypersurface, that $R^n \setminus X$ result into two connected regions, the outside D_0 and inside D_1 while D_1 itself is a compact manifold with boundary $\partial \overline{D_1} = X$.

The theorem implies that a surface enclosing volume in R^3 must separate inside and outside.

2. Theorem (Divergence theorem)¹⁵ Let $E \subset R^3$ be a compact region with boundary $S = \partial E$. For a vector field F .

$$\iiint_V \cdot F dV$$

The application of this theorem requires a consistent normal orientation on boundary surface.

3. Definition (Orientability)¹⁶

Let the manifold M be oriented in R^n . It is orientable if there exists a smooth map $n : M \rightarrow R^n$, where at each point we can define a corresponding continuous unit normal vector field $\vec{n}(p)$ on M and its size is $\|\vec{n}(p)\| = 1$.

A globally consistent normal vector field allows a surface to separate inside and outside.

4. Definition (Embedding)¹⁷

Let X and Y be two topological spaces, and $f : X \rightarrow Y$ is an injective continuous map. Let Z be the image of X in Y , a subspace of Y donated to the function $f' : X \rightarrow Z$ is bijective by restricting the range of f . We call that the map $f : X \rightarrow Y$ is topological embedding if f' is homeomorphism.

An embedding place a topological space in another larger space while preserves the its shape, connectivity and structure.

5. Definition (Immersion)^{3,18–20}

Let $f \rightarrow N$ be a smooth map. Then f is an immersion if its differential $df_{pM} \rightarrow T_{f(p)}N$ has rank equal to $\dim(M)$ for every point $p \in M$.

Immersion may contain self-intersections.

The Klein bottle is non-orientable.

Proposition. The Klein Bottle is non-orientable.

Proof. The Klein bottle can be represented as a quotient space obtained from the unit square $Q = [0, 1] \times [0, 1]$.^{21–23}




Figure. 3 The Klein bottle as a quotient space

with edge identifications:

$(0, y) \sim (1, y)$ and $(x, 0) \sim (x, 1)$ for all $x, y \in [0, 1]$.

The Klein bottle is defined as the quotient space $K = Q/\sim$.¹⁷

Suppose K is orientable. Thus, there must exist a continuous orientation on the tangent spaces $T_p K$ and a consistent oriented tangent basis on every point:

$$(V_{1(p)}, V_{2(p)})$$

For every tangent space $T_p K$ such that all transition maps keep the orientation.

Consider the identification map corresponding to the twisted edge:

$$\phi : [0, 1] \rightarrow [0, 1]$$

Defined by:

$$\phi(x) = 1 - x$$

The derivative of ϕ is

$$\frac{d\phi}{dx} = -1$$

Since $\det(d\phi) = -1 < 0$, the map ϕ reverses its orientation.

The oriented basis at a point P on the lower edge of Q :

$$(e_1, e_2)$$

becomes:

$$(-e_1, e_2)$$

After completing the loop, the orientation of the ordered basis reversed:

$$(e_1, e_2) \rightarrow (-e_1, e_2)$$

Thus, no globally consistent orientation can descend to the quotient space K . This contradicts the assumption that K is orientable. Hence K is non-orientable.

Embedding and Immersion

Proposition. The Klein bottle cannot be embedded in R^3 .

Proof. Assume the Klein bottle K can be embedded in R^3 . Since K is a compact closed embedded surface, the Jordan-Brouwer Separation Theorem states that $R^3 \setminus K$ has exactly two connected components: an inside and an outside. Thus, K would be the boundary separating those regions. The boundary of the compact three-dimensional region must have a consistent outward pointing normal unit vector, which means K is orientable. However, proposition 4.1 says that the Klein bottle is non-orientable, forming a contradiction. Therefore, the Klein bottle K cannot be embedded in R^3 .

Proposition. Klein Bottle is immersible in R^3 .

Proof. We must first prove that the Klein Bottle descends to a smooth map on the quotient space and then compute the partial derivatives to calculate the rank of the Jacobian Matrix to show the Klein Bottle is immersible in R^3 .

Step 1. Let the Klein Bottle be represented as a quotient of a square $[0, 2\pi] \times [0, 2\pi]$ with identifications:

$$(u, 0) \sim (u, 2\pi), (0, v) \sim (2\pi, 2\pi - v)$$

(Note: We represent the Klein Bottle as a unit square instead of a $[0, 1] \times [0, 1]$ unit square because the Klein Bottle immersion is built from rotations, where parametric equations will be involved and rotations are 2π periodic.)

Consider the standard figure-eight parametrization:

$$f(u, v) = \begin{pmatrix} (r + \cos(u/2) \sin(v) - \sin(u/2) \sin(2v)) \cos(u) \\ (r + \cos(u/2) \sin(v) - \sin(u/2) \sin(2v)) \sin(u) \\ \sin(u/2) \sin(v) + \cos(u/2) \sin(2v) \end{pmatrix}$$

And let:

$$x(u, v) = \left(r + \cos\left(\frac{u}{2}\right) \sin(v) - \sin\left(\frac{u}{2}\right) \sin(2v) \right) \cos(u)$$

$$y(u, v) = \left(r + \cos\left(\frac{u}{2}\right) \sin(v) - \sin\left(\frac{u}{2}\right) \sin(2v) \right) \sin(u)$$

$$z(u, v) = \sin\left(\frac{u}{2}\right) \sin(v) + 2 \cos\left(\frac{u}{2}\right) \sin(2v)$$

for $0 \leq u < 2\pi, 0 \leq v < 2\pi$ and $r > 2$




Figure. 4 The classical figure-eight immersion of the Klein bottle in R^3 .

For the horizontal edge identification $(u, 0) \sim (u, 2\pi)$ since,

$$\sin(0) = \sin(2\pi), \sin(2 \times 0) = \sin(4\pi) = 0$$

By substituting $v = 0$ or $v = 2\pi$ into the parametric equation:

$$x(u, 0) = r \cos(u) = x(u, 2\pi)$$

$$y(u, 0) = r \sin(u) = y(u, 2\pi)$$

$$z(u, 0) = 0 = z(u, 2\pi)$$

Where $f(u, 0) = f(u, 2\pi)$.

For vertical edge identification $(0, v) \sim (2\pi, 2\pi - v)$ first $f(0, v)$ is evaluated, by using

$$\cos\left(\frac{0}{2}\right) = 1, \cos(0) = 1, \sin\left(\frac{0}{2}\right), \sin(0) = 0$$

Yielding:

$$x(0, v) = r + \sin v$$

$$y(0, v) = 0$$

$$z(0, v) = \sin 2v$$

Second, $f(2\pi, 2\pi - v)$ is evaluated using:

$$\cos 2\pi = 1, \cos \pi = -1, \sin 2\pi = 0, \sin \pi = 0$$

And trigonometry identities:

$$\sin(2\pi - v) = -\sin v, \sin(2(2\pi - v)) = -\sin 2v$$

By substituting into parametric equations yield:

$$x(2\pi, 2\pi - v) = r + (-1)(-\sin v) - 0 = r + \sin v$$

$$y(2\pi, 2\pi - v) = 0$$

$$z(2\pi, 2\pi - v) = (-1)(-\sin 2v) = \sin 2v$$

Which is equivalent to the result for evaluating $f(0, v)$, therefore it can be concluded that $f(0, v) = f(2\pi, 2\pi - v)$.

Since both identifications satisfy:

$$f(u, 0) = f(u, 2\pi), f(0, v) = f(2\pi, 2\pi - v)$$

Hence, f descends to a defined smooth map on the quotient space.

Step 2. Now compute the rank of the Klein Bottle.

The following differential-geometric calculations use the standard parametrization and tangent vector frame-work for smooth surfaces.²⁴ The partial derivatives of $A(u, v)$ are $B(u, v)$ well computed.

Recall the map from the previous step:

$$f(u, v) = (x(u, v), y(u, v), z(u, v))$$

Let

$$A(u, v) = r + \cos\left(\frac{u}{2}\right) \sin(v) - \sin\left(\frac{u}{2}\right) \sin(2v)$$

$$B(u, v) = \sin\left(\frac{u}{2}\right) \sin(v) + 2 \cos\left(\frac{u}{2}\right) \sin(2v)$$

Were

$$x(u, v) = A(u, v) \cos u$$

$$y(u, v) = A(u, v) \sin u$$

$$z(u, v) = B(u, v)$$

With respect to u :

$$A_u = -\frac{1}{2} \sin \frac{u}{2} \sin v - \frac{1}{2} \cos \frac{u}{2} \sin 2v$$

$$B_u = \frac{1}{2} \cos \frac{u}{2} \sin v - \frac{1}{2} \sin \frac{u}{2} \sin 2v$$

With respect to v :

$$A_v = \cos \frac{u}{2} \cos v - 2 \sin \frac{u}{2} \cos 2v$$

$$B_v = \sin \frac{u}{2} \cos v + 2 \cos \frac{u}{2} \cos 2v$$

Step 3. compute $\frac{\partial f}{\partial u}$

Using the product rule results in it:

$$\frac{\partial x}{\partial u} = A_u \cos u - A \sin u$$

$$\frac{\partial y}{\partial u} = A_u \sin u + A \cos u$$

$$\frac{\partial z}{\partial u} = B_u$$

Thus,

$$\frac{\partial f}{\partial u} = \begin{pmatrix} A_u \cos u - A \sin u \\ A_u \sin u + A \cos u \\ B_u \end{pmatrix}$$

Step 4. Now computing $\frac{\partial f}{\partial v}$, again using the product rule:

$$\frac{\partial x}{\partial v} = A_v \cos u$$

$$\frac{\partial y}{\partial v} = A_v \sin u$$

$$\frac{\partial z}{\partial v} = B_v$$

Thus,

$$\frac{\partial f}{\partial v} = \begin{pmatrix} A_v \cos u \\ A_v \sin u \\ B_v \end{pmatrix}$$

Step 4. To compute $\frac{\partial f}{\partial v} \times \frac{\partial f}{\partial u}$ the cross-product matrix elements all determined:

$$\frac{\partial f}{\partial v} \times \frac{\partial f}{\partial u} = \begin{vmatrix} i & j & k \\ A_u \cos u - A \sin u & A_u \sin u + A \cos u & B_u \\ A_v \cos u & A_v \sin u & B_v \end{vmatrix}$$

$$\begin{aligned} &= i[(A_u \sin u + A \cos u) B_v - B_u (A_v \sin u)] \\ &\quad - j[(A_u \cos u - A \sin u) B_v - B_u (A_v \cos u)] \\ &\quad + k[(A_u \cos u - A \sin u) (A_v \sin u) - (A_u \sin u + A \cos u) (A_v \cos u)] \end{aligned}$$

then simplified:

$$= A_u B_v \sin u + A B_v \cos u - A_v B_u \sin u \\ - A_u B_v \cos u + A B_v \sin u + A_v B_u \cos u \\ - A A_v$$

and finally:

$$\frac{\partial f}{\partial v} \times \frac{\partial f}{\partial u} = \begin{pmatrix} A_u B_v \sin u + A B_v \cos u - A_v B_u \sin u \\ -A_u B_v \cos u + A B_v \sin u - A_v B_u \cos u \\ -A A_v \end{pmatrix}$$

Step 5. To prove that f defines an immersion, need to verify that

$$\text{rank}(df_{u,v}) = 2$$

For every point (u, v) .

Since the Jacobian matrix consists of the tangent vectors:

$$f_u = \frac{\partial f}{\partial u}, f_v = \frac{\partial f}{\partial v},$$

Rank two is equivalent to showing that these vectors are linearly independent.

A necessary and sufficient condition for linear independence is

$$f_u \times f_v \neq 0$$

Now compute the squared norm:

$$\|f_u \times f_v\|^2$$

Let

$$R_1(u, v) = A_u B_v \sin(u) + A B_v \cos(u) - A_v B_u \sin(u)$$

$$R_2(u, v) = -A_u B_v \cos(u) + A B_v \sin(u) - A_v B_u \cos(u)$$

$$R_3(u, v) = -A A_v$$

Which the squared norm of the cross-product equals:

$$\|f_u \times f_v\|^2 = R_1^2 + R_2^2 + R_3^2$$

Since each term is squared,

$$\|f_u \times f_v\|^2 \geq 0 \text{ occurs only if } R_1 = R_2 = R_3 = 0$$

While $R_3(u, v) = -A A_v$, vanishing requires either $A = 0$ or $A_v = 0$.

Substituting these cases into R_1 and R_2 shows that all three equations cannot hold simultaneously throughout the equation domain.

Thus:

$$\left\| \frac{\partial f}{\partial u} \times \frac{\partial f}{\partial v} \right\|^2 > 0 \text{ For all } (u, v) \in [0, 2\pi) \times [0, 2\pi)$$

Therefore, $\frac{\partial f}{\partial u}, \frac{\partial f}{\partial v}$, are linearly independent everywhere.

Hence

$$\text{rank}(df_{u,v}) = 2$$

For every point of the quotient domain. f is an immersion of the Klein bottle into R^3 .

Corollary. Every immersion of the Klein bottle into R^3 contains self-intersections.

Proof. Suppose an immersion of $K \rightarrow R^3$ has no self-intersections. Then the map would be injective. Since K is compact, an injective immersion would be an embedding. However, proposition 5.1 shows that K admits no embedding into R^3 . Hence every immersion must contain self-intersections.

Divergence Theorem

The Divergence theorem is another approach for volume enclosure. To apply the theorem, the manifold M must have a boundary in R^n . For an $n - 1$ -compact, oriented manifold M embedded in R^n , the theorem states that:

$$\text{thcanbe} \int_{\partial M} (\text{div } \mathbf{G}) dV = \int_M \langle \mathbf{G}, \mathbf{N} \rangle dV$$

Where G is a differentiable vector field, N is the consistently defined outward-pointing unit normal vector field.¹⁶ For the case of the Klein bottle, in R^3 without the boundary ∂M , the integral cannot be computed with an unidentified inside and outside; with in R^4 a continuous global unit normal field N is undefined fails to compute the volume using the Divergence theorem.

Discussion

This study shows that the Klein bottle cannot enclose a volume in R^3 because of its non-orientable and boundaryless properties. A surface that encloses a volume must have distinguishable inside and outside orientation, equivalent to being orientable. By the Jordan-Brouwer Separation theorem, an embedded compact space separating R^3 into an interior and exterior must be orientable, which the Klein bottle cannot be embedded into R^3 . However, it can be immersed in R^3 with self-intersections and eventually embedded in R^4 , by having an extra dimension to extend its shape. Non-orientability is a topological restriction for higher-dimensional topological figures, whereby a self-intersection is a unique shape of them in lower dimensions.

The analysis presents the relationship between topology and geometry, where the intrinsic topological properties can impose restrictions and affect how an object exists in Euclidean space. Furthermore, the paper clarifies the concepts of embeddings and immersions, which often confuse students. The use

of the Jordan-Brouwer separation theorem provides a topological explanation, while the use of the divergence theorem provides an additional geometric or vector-calculus perspective, helping students to connect and bring together topology and calculus. In academics, this paper combines several established concepts into one logical chain rather than discussing them separately in different chapters, as in textbooks. Its educational significance can be present in the second or third year of university, where students begin to learn topology and differential geometry that need them to cooperate with spatial structure and theorems; the comprehension of the known result is essential for them to do further studies.

The primary objective of the study was achieved: to provide an expository explanation to explain why the Klein bottle cannot enclose a volume in R^3 by establishing the non-orientability of the Klein bottle, explaining why the Klein bottle is non-orientable, which prevents volume generation. No new mathematical theorems or results were presented in the paper; therefore, success was achieved by clarity and synthesis of existing results. The study's conclusions were consistent with the original objectives.

Limitations

The paper is expository rather than original mathematical research and only includes established results rather than introducing a new theorem or proof. Thus, the findings are limited to existing mathematical knowledge. Advanced visualization tools or computational topology are not used in this paper. Conclusions strictly focus only on the Klein bottle and cannot be extended to all non-orientable manifolds, despite being in the same classification in topology.

Conclusion

Overall, the Klein bottle might appear to enclose a volume geometrically, but it is topologically impossible. Dimension and topology are the fundamental constraints of geometric objects, while volume enclosure associated with in-depth ideas regarding orientability, embedding, and the structure of the space. Topology helps to understand not only the shape but also the topological rules that control the way they exist in space.

Acknowledgment

I would like to thank Dr. Patricia Mitchler and Professor Derek Krepski for providing guidance and suggestions for the paper.

References

- 1 K. Polthier. Imaging maths Inside the Klein bottle. 1997.

- 2 Munkres James R. Topology (2nd edition). 1999.
- 3 Lee John M. Introduction to Smooth Manifolds. 2012.
- 4 Weeks Jeffrey Renwick. The Shape of Space. 1985.
- 5 Smith, A. "Piecewise linear immersions." Mathematical Proceedings of the Cambridge Philosophical Society, 68(1), 45–55, 1970. [<https://doi.org/10.1017/S0305004100001031>].
- 6 D. Cohen, L. Vandembroucq. Topological complexity of the klein bottle. Journal of Applied and Computational Topology. 2017.
- 7 H. H. Torriani. Profinite completions of the fundamental group of the klein bottle. Czechoslovak Mathematical Journal 35.4. pg. 511–514, 1985.
- 8 D. M. Davis. An n-dimensional klein bottle. Proceedings of the Royal Society of Edinburgh: Section A Mathematics. Vol. 149, pg. 1207–1221, 2019. <https://doi.org/10.1017/prm.2018.73>.
- 9 Cervone Davide P. Vertex-minimal simplicial immersions of the klein bottle in three-space. Geometriae Dedicata, 50. pg. 117–141, 1994.
- 10 Jakobson, D., Nadirashvili, N., & Polterovich, I. "Extremal Metric for the First Eigenvalue on a Klein Bottle." Canadian Journal of Mathematics, 58(2), 381–400, 2006. [<https://doi.org/10.4153/CJM-2006-016-0>].
- 11 J. Hirsch, E. Mäder-Baumdicker. A note on willmore minimizing klein bottles in euclidean space. Advances in Mathematics. Vol. 319, pg. 67–75, 2017. <https://doi.org/10.1016/J.AIM.2017.08.021>.
- 12 J. Eyll. Stability of systolic inequalities for the möbius strip and klein bottle. The Journal of Geometric Analysis 2025 35:11. Vol. 35, pg. 365–, 2025. <https://doi.org/10.1007/S12220-025-02197-9>.
- 13 B. Greene, D. Kabat, J. Levin, M. Porrtati. Klein bottle cosmology. 2026.
- 14 W. Schmaltz. THE JORDAN-BROUWER SEPARATION THEOREM. 2009.
- 15 Marsden Jerrold E., Tromba Anthony J. Vector Calculus. 1976.
- 16 Munkres James R. Analysis On Manifolds. CRC Press, 2018.
- 17 Lee John M. Introduction to Topological Manifolds. 2000.
- 18 Milnor John Willard. Topology from the Differentiable Viewpoint. 1965.
- 19 Guillemin Victor William, Pollack Alan. Differential Topology. 1974.
- 20 Hirsch Morris William. Differential Topology. 1976.
- 21 Massey William S. A Basic Course in Algebraic Topology. 1991.
- 22 Hatcher Allen. Algebraic Topology. 2001.
- 23 Armstrong Mark Anthony. Basic Topology. 1979.
- 24 do Carmo Manfredo Perdigão. Differential geometry of curves and surfaces. 1976.